



# Medway and Swale Integrated Care Partnership profile

Created by Medway Council Public Health Intelligence Team and Kent Public Health Observatory

*Transforming health and social care in Kent and Medway* is a partnership of all the NHS organisations in Kent and Medway, Kent County Council and Medway Council. We are working together to develop and deliver the Sustainability and Transformation Plan for our area.





## Summary part 1: Medway and Swale

Indicator	Compared to England
Life expectancy at birth (Male)	Similar
Life expectancy at birth (Female)	Similar
Smoking Prevalence in adults (18+) - current smokers (APS)	Similar
Percentage of adults (aged 18+) classified as overweight or obese	Worse
Children with excess weight Year 6, three year average	Similar
Percentage of physically inactive adults	Worse
Hospital stays for alcohol-related harm (Narrow definition)	Better
Air pollution: fine particulate matter	Not compared
Total number of prescribed antibiotic items per STAR-PU	Similar
Breast cancer screening coverage (females aged 50-70)	Better
Cervical cancer screening coverage (females aged 25-64)	Better
Bowel cancer screening coverage (persons aged 60-74)	Worse
Infant mortality rate	Similar
Low birth weight of term babies	Better
Stillbirth rate	Similar
Smoking status at time of delivery	Worse
Baby's first feed breastmilk	Better
AE attendances (0-4 years)	Better



## Summary part 2: Medway and Swale

Indicator	Compared to England
Percentage of 5 year olds with experience of visually obvious dental decay	Similar
Under 18s conception rate / 1,000	Worse
Emergency hospital admissions for asthma (< 19 yrs)	Similar
Emergency hospital admissions for epilepsy (< 19 yrs)	Worse
Emergency hospital admissions for diabetes (< 19 yrs)	Similar
Hospital admissions for mental health conditions (0-17 years)	Similar
Hospital admissions as a result of self-harm (10-24 years)	Better
Hospital admissions due to substance misuse (15-24 years)	Better
Hypertension: QOF prevalence (all ages)	Higher
Diabetes: QOF prevalence (17+)	Higher
CHD: QOF prevalence (all ages)	Lower
CKD: QOF prevalence (18+)	Lower
Stroke: QOF prevalence (all ages)	Lower
Deaths from circulatory disease, under 75 years	Similar
Deaths from all cancer, under 75 years	Worse
Cancer diagnosed at early stage (experimental statistics)	Not compared
Unplanned hospitalisation for chronic ACSC	Worse
Depression: Recorded prevalence (aged 18+)	Higher



## Summary part 3: Medway and Swale

Indicator	Compared to England
Serious Mental Illness: QOF prevalence (all ages)	Lower
Suicide rate (Persons)	Similar
Suicide rate (Male)	Worse
Estimated dementia diagnosis rate (aged 65 and over)	Not compared
Emergency hospital admissions due to falls (persons aged 65 and over)	Better
Emergency hospital admissions for hip fracture (persons aged 65 and over)	Similar
Osteoporosis: QOF prevalence (50+)	Lower




## Purpose



- Profiles have been created for each of the Integrated Care Partnerships (ICPs) in the Kent and Medway Integrated Care System (ICS).
- 1) Dartford, Gravesham and Swanley; 2) East Kent; 3) Medway and Swale; 4) West Kent.
- The aim of the profiles is to allow comparison between each of the ICPs and identify priority areas to focus work.



# Kent and Medway ICPs



Integrated Care Partnerships

-  Dartford, Gravesham and Swanley ICP
-  East Kent ICP
-  Medway and Swale ICP
-  West Kent ICP

Contains National Statistics data © Crown copyright and database right 2019  
Contains OS data © Crown copyright and database right 2019  
Produced by Medway Public Health Intelligence Team, Medway Council 2020-08-14




## Rationale

- The profiles contain five sections, which are based on key themes identified in the NHS Long Term Plan.
- 1) Demographics; 2) Prevention and Health Inequalities; 3) Best Start in Life; 4) Major Health Conditions; 5) Ageing Well.
- Key stakeholders were consulted to identify the indicators that should be included.
- Due to limitations in the available data, some indicators could not be included for all the priorities identified at this time.



## PCN hex map explained

- Some slides contain a hex map, which displays the indicator value at PCN level.
- Each hexagon represents a PCN, arranged according to its relative geography within the ICP.
- The PCN abbreviations used in the hex map are displayed below.







## Number grid explained

- Some slides contain a number grid, which displays the indicator value at different levels of geography.
- England -> Kent and Medway ICS -> ICP -> Small area.
- The small area type displayed depends on the data available for the indicator.
- Either District & Unitary Authority (UA) or Clinical Commissioning Group (CCG).

ICP value	England value
	ICS value
Small area values	Time period



## Small areas

- District & UA and CCG data have been mapped to the ICP boundaries as per the table below.
- Caveat: District & UA data does not align with the ICP areas exactly.
- For the purpose of this profile, data for whole District & UAs have been assigned to the ICP where the majority of residents reside.

Integrated Care Partnership	District & Unitary Authority	Clinical Commissioning Group
Dartford, Gravesham and Swanley	Dartford Gravesham	Dartford, Gravesham and Swanley CCG
East Kent	Ashford Canterbury Dover Folkestone & Hythe Thanet	Ashford CCG Canterbury and Coastal CCG South Kent Coast CCG Thanet CCG
Medway and Swale	Medway Swale	Medway CCG Swale CCG
West Kent	Maidstone Sevenoaks Tonbridge and Malling Tunbridge Wells	West Kent CCG



## Small areas

- Data at several small area levels has been used as building blocks to calculate the PCN values: Lower Super Output Area (LSOA), ward, general practice and school.
- LSOAs have a defined geographical boundary. On average the population is about 1,700 people so they can be thought of as representing a neighbourhood. There are 1,065 LSOAs within Kent and Medway.
- LSOAs and wards were assigned to PCNs on a first passed the post basis, e.g. LSOAs or wards were mapped to PCNs based on which PCN has the highest count of registered patients living in that LSOA/ward.
- School level data was assigned to PCNs based on the ward the school was located in. Only primary and nursery school data was used as this more likely reflects the child profile of the local area due to the larger catchment areas of secondary schools.



## ICP value and comparison

- The ICP values have been calculated from either LSOA, ward, general practice, school, District & UA or CCG level data using one of two methods:
- 1) Aggregated data: ICP values are created from aggregated counts and denominators, where data is available.
- 2) Small areas averaged: Where count and/or denominator data is not available, the ICP value is the median of the small area values.
- A RAG rating (red, amber, green) has been applied to the majority of indicators to show how well an area is performing compared to a benchmark (England). The RAG rating is assigned by comparing an area's value to a reference range, which was created using either confidence intervals (CIs) or a range around the England average (usually 5%). Green corresponds to a value that is better than England, red to a value that is worse, and amber indicates that there is no difference.
- Where it is inappropriate to label high or low values as 'better' or 'worse', for example osteoporosis prevalence, the terms 'higher' and 'lower' have been used with neutral colouring: shades of blue from light to dark. Such labelling does not imply that high values of these indicators, for example, are 'worse'.
- An indicator is shaded grey where it is inappropriate to apply a RAG rating due to the methods used in the calculation or the count is less than 10.

# DEMOGRAPHICS





# Population

## Age Profile for Medway and Swale

Total population: 402,657




- Male
- Female
- Kent & Medway
- England

Source: NHS Digital. Patients Registered at a GP Practice. 01JUL2020

## Projected population for MS

Percentage change from 2018



- 0-14
- 25-44
- 65-84
- 15-24
- 45-64
- 85+

Source: ONS. Population projections for local authorities. 2018 based.


Note: Population projections have been calculated by aggregating the local authority districts assigned to the ICP.



# Ethnicity


### School Census, January 2020 Nursery and Primary schools

- Medway and Swale
- Kent and Medway



### Census, 2011

- Medway and Swale
- Kent and Medway




Census source: NOMIS. 2011 Census. KS201EW)) - Ethnic group.  
 School Census source: GOV.UK. Department for Education. (2019). Schools, pupils and their characteristics: January 2020.



# Deprivation

National decile  
(1 = most deprived)

1 2 3 4 5 6 7 8 9 10



Contains National Statistics data © Crown copyright and database right 2019  
Contains OS data © Crown copyright and database right 2019  
Produced by Medway Public Health Intelligence Team, Medway Council 2020-08-14




# PREVENTION AND HEALTH INEQUALITIES






# Life expectancy at birth (Male)

PCNs in Medway and Swale. Compared to England:  
■ Better ■ Similar ■ Worse ■ Not compared



Trend data not available.




The rate in Medway and Swale is similar to England.

Value type: Years  
 Latest time period: 2013 - 17  
 Source: PHE, Fingertips, Indicator ID: 93283  
 ICP value calculation: Small areas averaged  
 Small area type: Ward to PCN  
 ICP RAG method: England plus/minus 2%




# Life expectancy at birth (Female)

PCNs in Medway and Swale. Compared to England:  
■ Better ■ Similar ■ Worse ■ Not compared



Trend data not available.





The rate in Medway and Swale is similar to England.

Value type: Years  
 Latest time period: 2013 - 17  
 Source: PHE, Fingertips, Indicator ID: 93283  
 ICP value calculation: Small areas averaged  
 Small area type: Ward to PCN  
 ICP RAG method: England plus/minus 2%




# Smoking Prevalence in adults (18+) - current smokers (APS)



The rate in Medway and Swale is similar to England.

Value type: Proportion - %  
 Latest time period: 2019  
 Source: PHE, Fingertips, Indicator ID: 92443  
 ICP value calculation: Small areas averaged  
 Small area type: District & UA (4/19-3/20)  
 ICP RAG method: England plus/minus 5%



There are data quality concerns with the figure for Dartford LA (see notes on the next slide).





## Smoking Prevalence in adults (18+) - current smokers (APS)

- Smoking prevalence is an estimate based on a sample of the population questioned in the Annual Population Survey run by the Office for National Statistics.
- The figure for Dartford LA in 2019 was high (26.4%) but with a very wide degree of uncertainty.
- This is most likely due to a small sample of people.
- Therefore the aggregate figure for DGS ICP should be interpreted with caution.




# Percentage of adults (aged 18+) classified as overweight or obese



The rate in Medway and Swale is worse than England.




Value type: Proportion - %  
 Latest time period: 2018/19  
 Source: PHE, Fingertips, Indicator ID: 93088  
 ICP value calculation: Small areas averaged  
 Small area type: District & UA (4/19-3/20)  
 ICP RAG method: England plus/minus 5%





# Children with excess weight Year 6, three year average

PCNs in Medway and Swale. Compared to England:  
■ Better ■ Similar ■ Worse ■ Not compared




The rate in Medway and Swale is similar to England.

Value type: Proportion - %  
 Latest time period: 2016/17 - 18/19  
 Source: PHE, Fingertips, Indicator ID: 93108  
 ICP value calculation: Aggregated data  
 Small area type: Ward to PCN  
 ICP RAG method: Confidence interval (95%) - Wilson Score method




# Percentage of physically inactive adults

<b>23</b>		<b>21</b>
		England
ICP: Medway and Swale		<b>21</b>
		Kent and Medway
<b>23</b>	<b>23</b>	2018/19
Medway	Swale	



The rate in Medway and Swale is worse than England.

Value type: Proportion - %  
 Latest time period: 2018/19  
 Source: PHE, Fingertips, Indicator ID: 93015  
 ICP value calculation: Small areas averaged  
 Small area type: District & UA (4/19-3/20)  
 ICP RAG method: England plus/minus 5%







# Hospital stays for alcohol-related harm (Narrow definition)


PCNs in Medway and Swale. Compared to England:

■ Better ■ Similar ■ Worse ■ Not compared



No trend data available.

Narrow definition: Admissions to hospital where the primary diagnosis is an alcohol-related condition, or a secondary diagnosis is an alcohol-related external cause.



The rate in Medway and Swale is better than England.

Value type: Indirectly standardised ratio per 100

Latest time period: 2013/14 - 17/18

Source: PHE, Fingertips, Indicator ID: 93240

ICP value calculation: Aggregated data


Small area type: Ward to PCN

ICP RAG method: Confidence interval (95%) - Byar's method




# Air pollution: fine particulate matter

10.6		8.9
		England
ICP: Medway and Swale		10.0
		Kent and Medway
11.2	10.0	2017
Medway	Swale	



Medway and Swale cannot be compared to England statistically.

Value type: Mean - µg/m3  
 Latest time period: 2017  
 Source: PHE, Fingertips, Indicator ID: 92924  
 ICP value calculation: Small areas averaged  
 Small area type: District & UA (4/19-3/20)  
 ICP RAG method: None applied





Last 8 years to 2017




# Total number of prescribed antibiotic items per STAR-PU

PCNs in Medway and Swale. Compared to England:  
■ Lower ■ Similar ■ Higher ■ Not compared



STAR-PU: Specific Therapeutic group Age-sex weightings Related Prescribing Unit  
The rate in Medway and Swale is similar to England.



Value type: Indirectly standardised ratio - per STAR-PU  
Latest time period: 2020 Q1  
Source: PHE, Fingertips, Indicator ID: 91900  
ICP value calculation: Small areas averaged  
Small area type: Practice to PCN  
ICP RAG method: England plus/minus 5%






# Breast cancer screening coverage (females aged 50-70)

PCNs in Medway and Swale. Compared to England:  
■ Better ■ Similar ■ Worse ■ Not compared



The rate in Medway and Swale is better than England.

Value type: Proportion - %  
 Latest time period: 2018/19  
 Source: PHE, Fingertips, Indicator ID: 91339  
 ICP value calculation: Aggregated data  
 Small area type: Practice to PCN  
 ICP RAG method: Confidence interval (99.8%) - Wilson Score method




Last 5 years to 2018/19

60  
40  
20  
0




# Cervical cancer screening coverage (females aged 25-64)

PCNs in Medway and Swale. Compared to England:  
 ■ Better ■ Similar ■ Worse ■ Not compared



The rate in Medway and Swale is better than England.



Value type: Proportion - %  
 Latest time period: 2018/19  
 Source: PHE, Fingertips, Indicator ID: 91341  
 ICP value calculation: Aggregated data  
 Small area type: Practice to PCN  
 ICP RAG method: Confidence interval (99.8%) - Wilson Score method






# Bowel cancer screening coverage (persons aged 60-74)

PCNs in Medway and Swale. Compared to England:  
■ Better ■ Similar ■ Worse ■ Not compared



The rate in Medway and Swale is worse than England.

Value type: Proportion - %  
 Latest time period: 2018/19  
 Source: PHE, Fingertips, Indicator ID: 92600  
 ICP value calculation: Aggregated data  
 Small area type: Practice to PCN  
 ICP RAG method: Confidence interval (99.8%) - Wilson Score method



Last 5 years to 2018/19




60  
40  
20  
0

# BEST START IN LIFE





# Infant mortality rate





The rate in Medway and Swale is similar to England.

Value type: Crude rate - per 1,000  
 Latest time period: 2016 - 18  
 Source: PHE, Fingertips, Indicator ID: 92196  
 ICP value calculation: Aggregated data  
 Small area type: District & UA (4/19-3/20)  
 ICP RAG method: Confidence interval (95%) - Byar's method






# Low birth weight of term babies





The rate in Medway and Swale is better than England.

Value type: Proportion - %  
 Latest time period: 2018  
 Source: PHE, Fingertips, Indicator ID: 20101  
 ICP value calculation: Aggregated data  
 Small area type: District & UA (4/19-3/20)  
 ICP RAG method: Confidence interval (95%) - Wilson Score method






# Stillbirth rate



Compared to England (---)

● Better ● Similar ● Worse ● Not compared



The rate in Medway and Swale is similar to England.

Value type: Crude rate - per 1,000

Latest time period: 2016 - 18

Source: PHE, Fingertips, Indicator ID: 92530



ICP value calculation: Aggregated data

Small area type: CCGs (2019/20)

ICP RAG method: Confidence interval (95%) - Byar's method




# Smoking status at time of delivery



The rate in Medway and Swale is worse than England.


Value type: Proportion - %  
 Latest time period: 2018/19  
 Source: PHE, Fingertips, Indicator ID: 93085  
 ICP value calculation: Aggregated data  
 Small area type: District & UA (4/19-3/20)  
 ICP RAG method: Confidence interval (95%) - Wilson Score method






# Baby's first feed breastmilk

<h1>70</h1>		<h1>67</h1>
		England
ICP: Medway and Swale		<h1>67</h1>
Kent and Medway		
<h1>71</h1>	<h1>65</h1>	2018/19
Medway	Swale	



Compared to England (---)

● Better ● Similar ● Worse ● Not compared



The rate in Medway and Swale is better than England.

Value type: Proportion - %

Latest time period: 2018/19

Source: PHE, Fingertips, Indicator ID: 93614

ICP value calculation: Aggregated data


Small area type: CCGs (2019/20)

ICP RAG method: Confidence interval (95%) - Wilson Score method



# AE attendances (0-4 years)

PCNs in Medway and Swale. Compared to England:  
■ Better ■ Similar ■ Worse ■ Not compared



The rate in Medway and Swale is better than England.

Value type: Crude rate per 1,000


Latest time period: 2018/19

Source: Hospital Episode Statistics (HES), NHS Digital

ICP value calculation: Aggregated data



Small area type: LSOA to PCN

ICP RAG method: Confidence interval (95%) - Byar's method






# Percentage of 5 year olds with experience of visually obvious dental decay



The rate in Medway and Swale is similar to England.

Value type: Proportion - %  
 Latest time period: 2019/20  
 Source: PHE, Fingertips, Indicator ID: 93563  
 ICP value calculation: Small areas averaged  
 Small area type: District & UA (pre 4/19)  
 ICP RAG method: England plus/minus 5%






# Under 18s conception rate / 1,000

<h1>23</h1>		<h1>17</h1>
		England
ICP: Medway and Swale		<h1>18</h1>
		Kent and Medway
<h1>23</h1>	<h1>24</h1>	2018
Medway	Swale	

Compared to England (---)  
 ● Better ● Similar ● Worse ● Not compared



The rate in Medway and Swale is worse than England.

Value type: Crude rate - per 1,000


Latest time period: 2018

Source: PHE, Fingertips, Indicator ID: 20401

ICP value calculation: Aggregated data

Small area type: District & UA (4/19-3/20)



ICP RAG method: Confidence interval (95%) - Byar's method





# Emergency hospital admissions for asthma (< 19 yrs)

PCNs in Medway and Swale. Compared to England:  
 ■ Better ■ Similar ■ Worse ■ Not compared



The rate in Medway and Swale is similar to England.

Value type: Crude rate - per 100,000


Latest time period: 2016/17 - 18/19

Source: Hospital Episode Statistics (HES), NHS Digital

ICP value calculation: Aggregated data

Small area type: LSOA to PCN

ICP RAG method: Confidence interval (95%) - Byar's method








# Emergency hospital admissions for epilepsy (< 19 yrs)

<h1 style="color: red;">112</h1> <p>ICP: Medway and Swale</p>		<h1 style="font-size: 2em;">76</h1> <p>England</p>
		<h1 style="color: yellow;">75</h1> <p>Kent and Medway</p>
<h1 style="color: red;">105</h1> <p>Medway</p>	<h1 style="color: red;">131</h1> <p>Swale</p>	<p>2018/19</p>

Compared to England (---)

- Better
- Similar
- Worse
- Not compared



The rate in Medway and Swale is worse than England.

Value type: Crude rate - per 100,000


Latest time period: 2018/19

Source: PHE, Fingertips, Indicator ID: 93142

ICP value calculation: Aggregated data


Small area type: CCGs (2019/20)

ICP RAG method: Confidence interval (95%) - Byar's method






# Emergency hospital admissions for diabetes (< 19 yrs)



Compared to England (---)

- Better
- Similar
- Worse
- Not compared



The rate in Medway and Swale is similar to England.

Value type: Crude rate - per 100,000


Latest time period: 2018/19

Source: PHE, Fingertips, Indicator ID: 93140

ICP value calculation: Aggregated data



Small area type: CCGs (2019/20)

ICP RAG method: Confidence interval (95%) - Byar's method






# Hospital admissions for mental health conditions (0-17 years)



The rate in Medway and Swale is similar to England.

Value type: Crude rate - per 100,000  
 Latest time period: 2018/19  
 Source: PHE, Fingertips, Indicator ID: 92761  
 ICP value calculation: Aggregated data  
 Small area type: CCGs (2019/20)  
 ICP RAG method: Confidence interval (95%) - Byar's method







# Hospital admissions as a result of self-harm (10-24 years)

PCNs in Medway and Swale. Compared to England:

■ Better ■ Similar ■ Worse ■ Not compared



The rate in Medway and Swale is better than England.

Value type: Directly standardised rate


Latest time period: 2018/19

Source: Hospital Episode Statistics (HES), NHS Digital

ICP value calculation: Aggregated data



Small area type: LSOA to PCN

ICP RAG method: Confidence interval (95%) - Dobson's method





# Hospital admissions due to substance misuse (15-24 years)



The rate in Medway and Swale is better than England.

Value type: Directly standardised rate - per 100,000


Latest time period: 2016/17 - 18/19

Source: PHE, Fingertips, Indicator ID: 92755

ICP value calculation: Small areas averaged

Small area type: CCGs (2019/20)

ICP RAG method: England plus/minus 5%





# MAJOR HEALTH CONDITIONS






# Hypertension: QOF prevalence (all ages)

PCNs in Medway and Swale. Compared to England:  
■ Lower ■ Similar ■ Higher ■ Not compared



The rate in Medway and Swale is higher than England.



Value type: Proportion - %  
 Latest time period: 2018/19  
 Source: PHE, Fingertips, Indicator ID: 219  
 ICP value calculation: Aggregated data  
 Small area type: Practice to PCN  
 ICP RAG method: Confidence interval (99.8%) - Wilson Score method






# Diabetes: QOF prevalence (17+)

PCNs in Medway and Swale. Compared to England:  
■ Lower ■ Similar ■ Higher ■ Not compared



The rate in Medway and Swale is higher than England.

Value type: Proportion - %  
 Latest time period: 2018/19  
 Source: PHE, Fingertips, Indicator ID: 241  
 ICP value calculation: Aggregated data  
 Small area type: Practice to PCN  
 ICP RAG method: Confidence interval (99.8%) - Wilson Score method



Last 5 years to 2018/19







# CHD: QOF prevalence (all ages)

PCNs in Medway and Swale. Compared to England:

■ Lower ■ Similar ■ Higher ■ Not compared



The rate in Medway and Swale is lower than England.

Value type: Proportion - %  
 Latest time period: 2018/19  
 Source: PHE, Fingertips, Indicator ID: 273  
 ICP value calculation: Aggregated data  
 Small area type: Practice to PCN  
 ICP RAG method: Confidence interval (99.8%) - Wilson Score method

● MS — England





Last 5 years to 2018/19




# CKD: QOF prevalence (18+)

PCNs in Medway and Swale. Compared to England:  
 ■ Lower ■ Similar ■ Higher ■ Not compared



The rate in Medway and Swale is lower than England.

Value type: Proportion - %  
 Latest time period: 2018/19  
 Source: PHE, Fingertips, Indicator ID: 258  
 ICP value calculation: Aggregated data  
 Small area type: Practice to PCN  
 ICP RAG method: Confidence interval (99.8%) - Wilson Score method





Last 5 years to 2018/19



# Stroke: QOF prevalence (all ages)

PCNs in Medway and Swale. Compared to England:  
 ■ Lower ■ Similar ■ Higher ■ Not compared



The rate in Medway and Swale is lower than England.

Value type: Proportion - %


Latest time period: 2018/19

Source: PHE, Fingertips, Indicator ID: 212

ICP value calculation: Aggregated data

Small area type: Practice to PCN

ICP RAG method: Confidence interval (99.8%) - Wilson Score method






## Deaths from circulatory disease, under 75 years

PCNs in Medway and Swale. Compared to England:

■ Better ■ Similar ■ Worse ■ Not compared



Trend data not available.



The rate in Medway and Swale is similar to England.

Value type: Indirectly standardised ratio per 100

Latest time period: 2013 - 17

Source: PHE, Fingertips, Indicator ID: 93256

ICP value calculation: Aggregated data

Small area type: Ward to PCN


ICP RAG method: Confidence interval (95%) - Byar's method




# Deaths from all cancer, under 75 years

PCNs in Medway and Swale. Compared to England:

■ Better 
 ■ Similar 
 ■ Worse 
 ■ Not compared



Trend data not available.



The rate in Medway and Swale is worse than England.

Value type: Indirectly standardised ratio per 100

Latest time period: 2013 - 17

Source: PHE, Fingertips, Indicator ID: 93254

ICP value calculation: Aggregated data


Small area type: Ward to PCN

ICP RAG method: Confidence interval (95%) - Byar's method




# Cancer diagnosed at early stage (experimental statistics)

53		52
		England
ICP: Medway and Swale		54
Kent and Medway		
49	59	2017
Medway	Swale	



Medway and Swale cannot be compared to England statistically.

Value type: Proportion - %  
 Latest time period: 2017  
 Source: PHE, Fingertips, Indicator ID: 90834  
 ICP value calculation: Aggregated data  
 Small area type: District & UA (4/19-3/20)  
 ICP RAG method: None applied







# Unplanned hospitalisation for chronic ACSC

PCNs in Medway and Swale. Compared to England:

■ Better ■ Similar ■ Worse ■ Not compared



The rate in Medway and Swale is worse than England.

Value type: Directly standardised rate per 100,000


Latest time period: 2018/19

Source: Hospital Episode Statistics (HES), NHS Digital

ICP value calculation: Aggregated data

Small area type: LSOA to PCN



ICP RAG method: Confidence interval (95%) - Dobson's method






# Depression: Recorded prevalence (aged 18+)

PCNs in Medway and Swale. Compared to England:  
■ Lower ■ Similar ■ Higher ■ Not compared



The rate in Medway and Swale is higher than England.

Value type: Proportion - %  
 Latest time period: 2018/19  
 Source: PHE, Fingertips, Indicator ID: 848  
 ICP value calculation: Aggregated data  
 Small area type: Practice to PCN  
 ICP RAG method: Confidence interval (99.8%) - Byar's method










# Serious Mental Illness: QOF prevalence (all ages)

PCNs in Medway and Swale. Compared to England:  
■ Lower ■ Similar ■ Higher ■ Not compared





The rate in Medway and Swale is lower than England.

Value type: Proportion - %  
 Latest time period: 2018/19  
 Source: PHE, Fingertips, Indicator ID: 90581  
 ICP value calculation: Aggregated data  
 Small area type: Practice to PCN  
 ICP RAG method: Confidence interval (99.8%) - Wilson Score method






# Suicide rate (Persons)



The rate in Medway and Swale is similar to England.

Value type: Directly standardised rate - per 100,000  
 Latest time period: 2016 - 18  
 Source: PHE, Fingertips, Indicator ID: 41001  
 ICP value calculation: Small areas averaged  
 Small area type: District & UA (4/19-3/20)  
 ICP RAG method: England plus/minus 5%






# Suicide rate (Male)

<h1>17</h1>		<h1>15</h1>
		England
ICP: Medway and Swale		<h1>15</h1>
		Kent and Medway
<h1>17</h1>	<h1>16</h1>	2016 - 18
Medway	Swale	

Compared to England (---)

- Better
- Similar
- Worse
- Not compared



The rate in Medway and Swale is worse than England.

Value type: Directly standardised rate - per 100,000


Latest time period: 2016 - 18

Source: PHE, Fingertips, Indicator ID: 41001

ICP value calculation: Small areas averaged

Small area type: District & UA (4/19-3/20)

ICP RAG method: England plus/minus 5%



Last 16 years to 2016 - 18


# AGEING WELL






# Estimated dementia diagnosis rate (aged 65 and over)

54		67
		England
ICP: Medway and Swale		60
Kent and Medway		
51	60	2020
Medway	Swale	



Medway and Swale cannot be compared to England statistically.

Value type: Proportion - %  
 Latest time period: 2020  
 Source: PHE, Fingertips, Indicator ID: 92949  
 ICP value calculation: Aggregated data  
 Small area type: District & UA (4/19-3/20)  
 ICP RAG method: None applied



Last 4 years to 2020




# Emergency hospital admissions due to falls (persons aged 65 and over)

<h1 style="color: green;">1778</h1>		<h2 style="color: black;">2198</h2> <p>England</p>
		<h2 style="color: orange;">2157</h2> <p>Kent and Medway</p>
ICP: Medway and Swale		
<h2 style="color: green;">1768</h2> <p>Medway</p>	<h2 style="color: green;">1788</h2> <p>Swale</p>	<p>2018/19</p>

Compared to England (- - -)

● Better ● Similar ● Worse ● Not compared



The rate in Medway and Swale is better than England.

Value type: Directly standardised rate - per 100,000


Latest time period: 2018/19

Source: PHE, Fingertips, Indicator ID: 22401

ICP value calculation: Small areas averaged

Small area type: District & UA (4/19-3/20)

ICP RAG method: England plus/minus 5%




Last 9 years to 2018/19




# Emergency hospital admissions for hip fracture (persons aged 65 and over)

PCNs in Medway and Swale. Compared to England:

■ Better ■ Similar ■ Worse ■ Not compared



Trend data not available.



The rate in Medway and Swale is similar to England.

Value type: Indirectly standardised ratio per 100

Latest time period: 2013/14 - 17/18

Source: PHE, Fingertips, Indicator ID: 93241

ICP value calculation: Aggregated data



Small area type: Ward to PCN

ICP RAG method: Confidence interval (95%) - Byar's method



# Osteoporosis: QOF prevalence (50+)

PCNs in Medway and Swale. Compared to England:  
■ Lower ■ Similar ■ Higher ■ Not compared



The rate in Medway and Swale is lower than England.

Value type: Crude rate - %  
 Latest time period: 2018/19  
 Source: PHE, Fingertips, Indicator ID: 90443  
 ICP value calculation: Aggregated data  
 Small area type: Practice to PCN  
 ICP RAG method: Confidence interval (99.8%) - Wilson Score method

